



## Basal cortisol's relation to testosterone changes may not be driven by social challenges



Keith M. Welker<sup>a,\*</sup>, Smrithi Prasad<sup>b</sup>, Sanjay Srivastava<sup>b</sup>, Pranjal H. Mehta<sup>b</sup>

<sup>a</sup> Department of Psychology, University of Massachusetts Boston, United States of America

<sup>b</sup> Department of Psychology, University of Oregon, United States of America

### ARTICLE INFO

#### Keywords:

Testosterone  
Cortisol  
Competition  
Stress  
Cross-talk

### ABSTRACT

Multiple studies show a negative correlation between basal cortisol and testosterone changes in the presence of competition and social-evaluative stressors. These negative associations are proposed to be derived from psychological responses to competition and social-evaluative stress. However, we argue that the association between basal cortisol and testosterone change may instead be a statistical consequence of positively associated variables. In this paper, we present a mathematical rationale for this alternative explanation and examples from two studies that are consistent with this alternative explanation. Both studies show that the associations between basal cortisol and testosterone change have covariance patterns consistent with this alternative possibility. We conclude that the often-found positive association between basal cortisol and basal testosterone opens the door for alternative explanations of the basal cortisol-testosterone change association rooted in the patterns of associations between hormones measured over time. We also suggest future research directions and methods for testing alternative explanations.

The functional “cross-talk” (e.g., [Viau, 2002](#)) between the hypothalamic-pituitary-gonadal (HPG) and hypothalamic-pituitary-adrenal (HPA) axes is an increasingly studied area of research in the field of psychoneuroendocrinology. Together, testosterone and cortisol may jointly regulate social behaviors and traits (see [Mehta and Prasad, 2015](#) for a review). Additionally, simultaneous, coupled elevations of testosterone and cortisol characterize psychopathology and responses to stress (e.g., [Marceau et al., 2015](#); [Shirtcliff et al., 2015](#)). Overall, research on cross-talk and interactions between these axes is being increasingly used to understand social behaviors and personality traits.

A recent line of research investigating HPA-HPG cross-talk has examined correlations between basal cortisol and changes in testosterone from before to after a competition or social-evaluative stressor. Testosterone often increases in the presence of competition or when status is challenged ([Archer, 2006](#); [Casto and Edwards, 2016](#); [Geniole et al., 2017](#)). These testosterone increases tend to facilitate dominant behavioral responses in the face of social challenges ([Wingfield et al., 1990](#)). Moreover, because of evidence for cross-talk between the HPA and HPG axes, researchers have investigated whether basal cortisol is associated with the observed testosterone changes during social challenges. [Bedgood and colleagues](#) demonstrated a negative correlation between basal cortisol and testosterone reactivity during a socially evaluative stressor ([Bedgood et al., 2014](#)). Specifically, lower basal

cortisol levels predicted increased testosterone reactivity to the stressor. Additionally, [Edwards and Casto \(2015\)](#) extended these results by reporting a similar negative relation between basal cortisol and testosterone reactivity during athletic competitions. These findings are parallel to those of [Mehta and Josephs \(2006\)](#) and [Maestripieri et al. \(2010\)](#), that also show moderate negative associations between basal cortisol and testosterone changes in response to laboratory competitions and psychosocial stressors.

Based on these results, an elegant conclusion is that the association between basal cortisol and testosterone reactivity during competitions or stressors corresponds to psychological states and behaviors in stressful and competitive contexts ([Bedgood et al., 2014; 2015](#)). In particular, [Bedgood et al. \(2014\)](#) proposed that individuals with high basal cortisol levels perceive social evaluative situations as a threat to status rather than as a dominance challenge, leading to a blunted testosterone response ([Bedgood et al., 2014](#)). A similar psychological mechanism during athletic competition may explain the negative association between basal cortisol and testosterone change in the [Edwards and Casto \(2015\)](#) study. For brevity in this article, we refer to this potential explanation as the “social challenge explanation” of the association between basal cortisol and testosterone changes. The social challenge explanation claims that the negative association between basal cortisol and testosterone reactivity is a function of HPA-HPG

\* Corresponding author.

E-mail address: [Keith.Welker@umb.edu](mailto:Keith.Welker@umb.edu) (K.M. Welker).

cross-talk in social-evaluative stress and competitive contexts, particularly where status is contested.

This compelling social challenge explanation of the basal cortisol-testosterone change correlation would indeed be influential in moving psycho-neuroendocrine theory of competition and status forward. However, we argue it is premature to conclude that this correlation is produced by psychological factors. We propose another possibility for the association between basal cortisol and testosterone changes. A critical step in establishing the veracity of the social challenge explanation involves ruling out potential alternative explanations. We propose that the association between basal cortisol and testosterone changes is a direct consequence of covariance patterns between hormones measured over time and provide empirical evidence consistent with this alternative explanation. At this point, it is unknown what factors may create these covariance patterns, and if the factors are social psychological or physiological in nature. Therefore, it is necessary to explain what factors lead a negative basal cortisol and testosterone change association when measured over time. Although we refer to this as an “alternative possibility” or “explanation”, there may be many other possibilities that enable the conditions we identify contributing to a negative basal cortisol-testosterone change association. In this paper, we describe a rationale for this alternative statistical possibility and present data that are consistent with this alternative explanation. If this possibility is true, a critical next step is to determine what factors produce the covariance patterns that lead to a negative correlation between basal cortisol and testosterone changes. To do so, we also provide researchers with directions moving forward to investigate this alternative possibility.

## 1. The alternative statistical possibility

We propose that the negative association between basal cortisol and testosterone reactivity may be a statistical consequence of positively associated variables (i.e., basal testosterone and basal cortisol concentrations; Mehta and Josephs, 2010; Mehta et al., 2015a, 2015b, 2015c; Popma et al., 2007; Welker et al., 2016). More specifically, this association could be a consequence of the mathematics behind a correlation between two variables—in this case, testosterone and cortisol—measured at different times. Here, we present the argument that a negative covariance (and also a correlation) between basal cortisol and testosterone changes (assessed as difference scores) is inevitable, provided that the association between basal cortisol and basal testosterone (i.e. cortisol and testosterone measured at the same time) is stronger than the association between basal cortisol and testosterone measured at a later point. To illustrate this, we draw from the equation of a covariance between two variables, denoted here as X and Y in Eq. (1):

$$\text{cov}(Y, X) = \frac{\sum_{i=1}^n (Y_i - \bar{Y})(X_i - \bar{X})}{n - 1} \quad (1)$$

In this equation, we substitute basal cortisol and testosterone changes into the values of Y and X, respectively. Eq. (2) presents an algebraic statement of the covariance between basal cortisol (C1) and the change from basal testosterone (T1) to a later measure of testosterone (T2) as ( $\text{cov}(C1, [T2 - T1])$ ). From this, we derive (Eq. (2)) that the covariance of basal cortisol (C1) and testosterone changes from baseline (T2 - T1) is equal to the covariance of basal cortisol and post-competition testosterone ( $\text{cov}(C1, T2)$ ) minus the covariance of basal testosterone and cortisol ( $\text{cov}(C1, T1)$ ).

$$\begin{aligned} \text{cov}(C1, [T2 - T1]) &= \frac{\sum_{i=1}^n (C1_i - \bar{C1})[(T2_i - T1_i) - (\bar{T2} - \bar{T1})]}{n - 1} \\ &= \frac{\sum_{i=1}^n (C1_i - \bar{C1})(T2_i - \bar{T2}) - (T1_i - \bar{T1})}{n - 1} \end{aligned} \quad (2)$$

$$\begin{aligned} &= \frac{\sum_{i=1}^n [(C1_i - \bar{C1})(T2_i - \bar{T2}) - (C1_i - \bar{C1})(T1_i - \bar{T1})]}{n - 1} \\ &= \frac{\sum_{i=1}^n [(C1_i - \bar{C1})(T2_i - \bar{T2})]}{n - 1} - \frac{\sum_{i=1}^n [(C1_i - \bar{C1})(T1_i - \bar{T1})]}{n - 1} \\ &= \text{cov}(C1, T2) - \text{cov}(C1, T1) \end{aligned}$$

Based on this equation, if the covariance between basal testosterone and basal cortisol ( $\text{cov}(C1, T1)$ ) is greater than the covariance between basal cortisol and post-competition testosterone ( $\text{cov}(C1, T2)$ ), it follows that a negative covariance between testosterone changes and basal cortisol will occur. This is because this pattern of covariances will result in a negative value for  $\text{cov}(C1, [T2 - T1])$ . More simply, subtracting a larger positive quantity from a smaller positive quantity provides a negative quantity.

A correlation is mathematically linked to a covariance, as bivariate correlation coefficients in the form of Pearson's  $r$  are computed by dividing a covariance ( $\text{cov}(Y, X)$ ) by the product of the standard deviations ( $\sigma_Y$  and  $\sigma_X$ ) of both variables (Eq. (3)).

$$r = \frac{\text{cov}(Y, X)}{\sigma_Y \sigma_X} \quad (3)$$

Indeed, statistically, a correlation can be interpreted as a “standardized covariance” (Rodgers and Nicewander, 1988). Therefore, this potential statistical explanation will influence correlational analysis in the same way provided that the variances are not changing over time.<sup>1,2</sup>

## 2. Evidence consistent with the statistical explanation

According to the social challenge explanation, competition or stressors lead to a negative association between basal cortisol and testosterone reactivity. However, the alternative statistical possibility argues that the negative association between basal cortisol and testosterone reactivity should emerge across a variety of contexts, provided the condition basal testosterone and basal cortisol are positively associated and simultaneously measured hormones are more strongly correlated than measures of hormones from different times. Although the circumstances arising to this condition need to be explained by future research, basal testosterone and basal cortisol have a well-established positive correlation from previous literature (e.g., Mehta and Josephs, 2010; Mehta et al., 2015a, 2015b, 2015c; Popma et al., 2007; Welker et al., 2016; Zilioli and Watson, 2013). Furthermore, simultaneously measured variables that are correlated across time often have stronger synchronous correlations compared to when multiple measurements of variables are correlated across time in lagged correlations (Kenny, 1975; Kenny and Campbell, 1989), which is often the case with hormones (e.g., Liening et al., 2010; Welker et al., 2017). Indeed, many cross-lagged analyses show this pattern of correlations, where synchronous correlations between two variables are stronger than lagged correlations (e.g., Racine et al., 2016; Roest et al., 2016; Spurk and Abele, 2014; Wirtz et al., 2014). Therefore, if testosterone and cortisol have a sizeable synchronous correlation and the lagged correlations between testosterone and cortisol are weaker than the synchronous correlation, then the relationship between basal cortisol and testosterone change will be negative.

To demonstrate this alternative possibility, we analyzed archival data to test these associations across social contexts. To show data consistent with the alternative statistical possibility, we examined the patterns of covariance between basal cortisol and testosterone change

<sup>1</sup> Bedgood et al. (2014) and Edwards and Casto (2015) do not report correlations between individual testosterone and cortisol samples in either paper. Thus, it is not possible to compare correlations between simultaneously sampled testosterone and cortisol to cortisol and testosterone sampled at different times.

<sup>2</sup> For a related statistical explanation based on regression to the mean, see Maestripieri et al. (2010).

to determine whether the data conform to Eq. (2) mentioned above. Investigating both correlations and covariances, we also examined whether the association between testosterone and cortisol concentrations measured at the same time (i.e.,  $cov(C1, T1)$ ,  $cov(C2, T2)$ ) are stronger than those of the testosterone and cortisol measured at different times (i.e.,  $cov(C1, T2)$ ,  $cov(C2, T1)$ ).

In a recently published paper (Mehta et al., 2015a), we measured testosterone and cortisol reactivity during a face-to-face negotiation task (Study 1) and across multiple rounds of the Ultimatum Game (Study 2). In Study 1, participants ( $N = 70$ , 39% Women) provided a baseline saliva sample, were randomly paired with a partner, and assigned the role of a buyer or seller. Participants then engaged in a negotiation task involving selling a hypothetical pharmaceutical plant (the Synertech-Dosagen case; Galinsky and Mussweiler, 2001). Participants negotiated for approximately 15 min and provided a second saliva sample 20 min after the negotiation. In Study 2, participants ( $N = 115$ , 53% Women) provided a saliva sample and then played 30 one-shot rounds of the Ultimatum Game (Güth et al., 1982) for approximately 10 min. The Ultimatum game is an economic decision-making paradigm assessing whether people will retaliate against unfair treatment. In the game, a participant can pay a financial cost to punish the other participant for making an unfair offer. After the game, participants provided a second saliva sample to assess testosterone and cortisol roughly 20 min after the completion of the Ultimatum Game.

We took several measures to calculate changes in hormones. Cortisol values were log transformed to correct for skewness before this difference was calculated and then standardized, whereas testosterone was standardized within men and women separately. Absolute change in testosterone and cortisol were calculated by subtracting transformed Time 2 hormone concentrations from transformed Time 1 hormone concentrations (after the previously mentioned transformations were applied; See Mehta et al., 2015a).

Correlations in both studies between testosterone and cortisol levels at the corresponding time points, as well as testosterone and cortisol change, are presented in Table 1. To demonstrate the alternative statistical possibility, we examined the difference between the covariance between testosterone and cortisol at Time 1 and the covariance between Time 1 cortisol and post-negotiation testosterone (measured at Time 2) for both studies. Below, we present the covariances between Time 1 cortisol and testosterone changes ( $\sigma(C1, [T2 - T1])$ ) along with the covariance between Time 1 cortisol and testosterone ( $\sigma(C1, T1)$ ) and Time 1 cortisol and time 2 testosterone ( $\sigma(C1, T2)$ ). Cases were excluded listwise so that covariances represented only those with no missing data for C1, T1, and T2. These covariances also illustrate the principle that  $\sigma(C1, [T2 - T1])$  is derived from  $\sigma(C1, T2) - \sigma(C1, T1)$ :

$$\begin{aligned} \text{Study 1: } \sigma(C1, [T2 - T1]) &= \sigma(C1, T2) - \sigma(C1, T1) \\ &= 0.06 - 0.21 = -0.15 \end{aligned}$$

$$\begin{aligned} \text{Study 2: } \sigma(C1, [T2 - T1]) &= \sigma(C1, T2) - \sigma(C1, T1) \\ &= 0.14 - 0.30 = -0.16 \end{aligned}$$

Consistent with the alternative statistical possibility, these results show that in both studies that  $\sigma(C1, T1)$  is indeed greater than  $\sigma(C1, T2)$ , which results in a negative  $\sigma(C1, [T2 - T1])$ .

To supplement this primary evidence for our alternative explanation, we also examined correlations, excluding missing cases listwise. Expectedly, the pattern of correlations was similar to pattern of covariances. Similar to the findings of Bedgood et al. (2014) and Edwards and Casto (2015), basal cortisol was negatively associated with testosterone change in Study 1 ( $r(67) = -0.24$ ,  $p = 0.047$ ) and Study 2 ( $r(109) = -0.26$ ,  $p = 0.007$ ). Additionally, the basal testosterone and basal cortisol correlations ( $r(67) = 0.23$ ,  $p = 0.057$  and  $r(109) = 0.30$ ,  $p = 0.002$  for Studies 1 and 2, respectively) were generally of more robust magnitude than testosterone and cortisol correlations from measures at different times ( $r$ s from 0.04 to 0.24, See Table 1). For interested readers, we also examined the pattern of correlations and covariances separately for men and women, noting that they are of

similar magnitude as the pattern of findings across men and women (see Tables S1 and S2 in the Online Supplemental Materials). Furthermore, since basal testosterone was correlated with testosterone changes in Studies 1 ( $r(67) = -0.30$ ,  $p = 0.016$ ) and 2 ( $r(109) = -0.32$ ,  $p = 0.001$ ), we also controlled for basal testosterone when examining the basal cortisol and testosterone change association. Doing so resulted in a negative, albeit nonsignificant, partial correlation in Study 1 ( $r_p(67) = -0.19$ ,  $p = 0.131$ ) and a marginally-significant negative correlation in Study 2 ( $r_p(109) = -0.18$ ,  $p = 0.058$ ). Although the differences in these correlations was nonsignificant ( $ps \geq 0.276$ ) and our data was likely under-powered to detect such a change, this relationship was slightly attenuated but still in the same negative direction, when accounting for both basal hormone measurements.

We note that the association between post-interaction testosterone and cortisol was diminished ( $r(67) = -0.02$ ,  $p = 0.855$  and  $r(109) = 0.15$ ,  $p = 0.120$  for Studies 1 and 2, respectively). This reduced association between post-task testosterone and cortisol may exist because post-task testosterone and cortisol are influenced by more situational and individual difference factors in addition to the interaction. For instance, a growing body of work suggests several factors that may promote hormonal reactivity to competitive and stressful interactions, including enjoyment of the activity (Mehta et al., 2015b), competitive outcomes (e.g., Carré et al., 2013; Mehta et al., 2015b), and aggressiveness (e.g., Carré et al., 2010; Carré et al., 2013). These factors may have contributed to more unexplained variation in post-interaction hormones, relative to basal hormones. We return to this point later.

### 3. Discussion

Overall, our calculations and the patterns in our data suggest that the negative association between basal cortisol and testosterone changes may be a statistical consequence of positive correlations between basal cortisol and basal testosterone. Therefore, it is plausible that observations of HPA-HPG cross-talk (i.e., associations between basal cortisol and testosterone change) may not be relevant to adapting to social challenges. With the goal of moving theory forward, we propose several future directions to further establish whether the social challenge explanation may be true. We also speculate about the possible reasons that may give rise to the covariance patterns between hormones measured at different times that lead to the negative correlation between basal cortisol and testosterone changes.

First, a critical step forward to test the social challenge explanation is to experimentally manipulate the presence of social challenges, such as randomly assigning participants to a stressful/competitive condition or a control condition. Although concerns over this limitation have been raised (Bedgood et al., 2014), no research, including data presented from the present research, has yet compared the relationship between basal cortisol and testosterone reactivity in the presence and absence of social challenges. Such a study might use moderated regression analysis to test for a significant interaction between basal cortisol and a variable coding for the experimental condition predicting testosterone responses. This approach would be superior to examining correlations between basal cortisol and testosterone changes separately within different experimental conditions (for an example of the latter approach, see Mehta and Josephs, 2006). Indeed, it would provide a proper test of whether the experimentally manipulated social context significantly changes (that is, moderates) the relationship between basal cortisol and testosterone responses.<sup>3</sup> If the social challenge explanation is true, then researchers will find significantly more pronounced negative associations between basal cortisol and testosterone reactivity in

<sup>3</sup> Both Bedgood et al. (2014) and Edwards and Casto (2015) used the terms “moderator” or “moderates” in their discussion of social contexts without sufficient evidence. We caution against the use of these terms in study designs that are not designed to test for moderation. Instead, we recommend terms such as association or relationship for correlational studies like Bedgood et al. (2014) and Edwards and Casto (2015).

**Table 1**  
Associations between testosterone, cortisol, and hormonal changes in Mehta et al. (2015a).

Correlations							
Study 1	1	2	3	4	5	6	
1. Basal Cortisol (lg10, standardized)	–						
2. Basal Testosterone (standardized within sex)	0.23 <sup>†</sup>	–					
3. Post-interaction Cortisol (lg10, standardized)	0.53 <sup>***</sup>	0.04	–				
4. Post-interaction Testosterone (standardized within sex)	0.06	0.77 <sup>***</sup>	–0.02	–			
5. Cortisol Change	–0.60 <sup>***</sup>	–0.22 <sup>*</sup>	0.37 <sup>**</sup>	–0.09	–		
6. Testosterone Change	–0.24 <sup>†</sup>	–0.30 <sup>†</sup>	–0.09	0.39 <sup>***</sup>	0.18	–	
Study 2	1	2	3	4	5	6	
1. Basal Cortisol (lg10, standardized)	–						
2. Basal Testosterone (standardized within sex)	0.30 <sup>**</sup>	–					
3. Post-interaction Cortisol (lg10, standardized)	0.76 <sup>***</sup>	0.24 <sup>†</sup>	–				
4. Post-interaction Testosterone (standardized within sex)	0.14	0.81 <sup>***</sup>	0.15	–			
5. Cortisol Change	–0.37 <sup>***</sup>	–0.10	0.32 <sup>***</sup>	0.01	–		
6. Testosterone Change	–0.26 <sup>**</sup>	–0.32 <sup>***</sup>	–0.14	0.30 <sup>***</sup>	0.18 <sup>†</sup>	–	
Covariances							
Study 1	1	2	3	4	5	6	
1. Basal Cortisol (lg10, standardized)	0.91						
2. Basal Testosterone (standardized within sex)	0.21	0.88					
3. Post-interaction Cortisol (lg10, standardized)	0.42	0.03	0.68				
4. Post-interaction Testosterone (standardized within sex)	0.06	0.70	–0.02	0.95			
5. Cortisol Change	–0.50	–0.18	0.26	–0.08	0.76		
6. Testosterone Change	–0.15	–0.18	–0.05	0.25	0.10	0.43	
Study 2	1	2	3	4	5	6	
1. Basal Cortisol (lg10, standardized)	1.02						
2. Basal Testosterone (standardized within sex)	0.30	1.02					
3. Post-interaction Cortisol (lg10, standardized)	0.76	0.24	0.97				
4. Post-interaction Testosterone (standardized within sex)	0.14	0.82	0.15	1.01			
5. Cortisol Change	–0.26	–0.07	0.21	0.01	0.47		
6. Testosterone Change	–0.16	–0.20	–0.09	0.19	0.08	0.39	

Outliers in Study 1 were Winsorized to 3 SDs after transformations were applied. There were no outliers identified in Study 2. Covariances and correlations are presented excluding cases listwise. Because the standardized variables had outliers Winsorized after standardizing the variables, the diagonal covariances are not precisely equal to 1.

Note:

- <sup>†</sup>  $p < 0.10$ .
- <sup>\*</sup>  $p < 0.05$ .
- <sup>\*\*</sup>  $p < 0.01$ .
- <sup>\*\*\*</sup>  $p < 0.001$ .

competition or stressful situations compared to control conditions (as Bedgood et al., 2014 suggest). Future research with this approach will help determine if the presence of a social challenge or competition alters the association between basal cortisol and testosterone change. However, if the social challenge explanation were not true, this association would be constant regardless of the psychological or social context. It is also important that research testing this explanation be fueled by studies with larger sample sizes, as the empirical data presented in this paper and by previous researchers (Bedgood et al., 2014; Edwards and Casto, 2015) are likely underpowered for detecting statistical moderation. Future work with larger sample sizes will provide adequate statistical power to investigate whether social contexts and challenges systematically alter how basal cortisol and testosterone changes are associated.

By experimentally manipulating the social context, researchers might be able to parse out the psychologically-meaningful variability in the basal cortisol and testosterone change relationship from the variability that is a by-product of statistical covariances. It is possible that by controlling for basal hormones in the analyses of hormone changes, either through partial correlations, regression models, or lagged analyses using repeated measures (e.g., multilevel modeling, lagged path analysis), researchers might estimate variability in hormone changes

independent of the influence of basal hormones. Other researchers have used residualized values of change to index hormonal changes created from regressing a later measure of hormones on an earlier or basal hormone measurement (e.g., Smith and Apicella, 2017; Welker et al., 2017). However, this approach may not fully resolve this problem. In the current research, doing this resulted in smaller negative partial correlations between basal cortisol and testosterone changes, suggesting that basal testosterone accounts for some of the variability in this relationship. However, controlling for these factors did not substantially or significantly decrease the negative association between testosterone changes and basal cortisol. Although these approaches may not fully mitigate the statistical explanation we raise, they will aid in examining hormone changes independent of the influence basal hormones.

Another direction is for researchers to share the correlations between basal hormones and hormone changes in future publications, or from previously collected data through data sharing initiatives such as the open science framework (<https://osf.io/>), when possible. As research on HPG-and HPA cross-talk unfolds, researchers might begin investigating meta-analytic associations between basal hormones and changes. The availability of this data will help researchers compare studies presenting basal and hormone change correlations across a

variety of social and experimental contexts. Moreover, meta-analytically analyzing the differences in basal and hormone change correlations across different contexts would also help determine the extent to which specific contextual factors moderate the association between basal cortisol and testosterone changes.

This paper helps demonstrate why a negative correlation between basal cortisol and testosterone changes would occur given the correlations between two hormones measured at baseline compared to lagged correlations. However, a larger remaining issue needs to be addressed: Why is the covariance between basal cortisol and basal testosterone greater than the covariance between basal cortisol and a lagged measure of testosterone? We speculate that many of the laboratory methods that researchers employ when assessing hormonal changes (e.g., stress-induction tasks, competitive interactions) may induce a variety of differential hormone responses depending on the person. This may lead to reduced correlations between hormones measured at different times or after the laboratory task, relative to the baseline hormones. Indeed, our data followed this pattern, with correlations being the strongest between single measures of testosterone and cortisol at baseline. However, by comparing data from many different research approaches and hormonal changes measured in the absence of a psychologically-relevant task, researchers will have a better sense of why correlations decrease when using lagged hormone measures compared to baseline measures. Furthermore, researchers will be able to detect whether the negative correlation between basal cortisol and testosterone changes occurs across many different social contexts.

Finally, an additional next step is to employ pharmacological manipulations of cortisol (e.g., Metyrapone or Dexamethasone) to examine whether cortisol can causally alter testosterone reactivity. Such a design could test whether exogenously-administered cortisol diminishes testosterone reactivity to social challenges, supporting the presence of social challenge-dependent HPA-HPG crosstalk. These new directions in research may shed new light on HPA and HPG cross-talk in the presence of social challenges, and its potential influence on behavior.

### Conflict of interest

None of the authors report any conflict of interest.

### Acknowledgements

This research was supported by a National Science Foundation grant awarded to PHM (Award Number: 1451848). This funding sponsor was not involved in study design; in the collection, analysis and interpretation of data; in the writing of the report; or in the decision to submit the article for publication.

### Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.psyneuen.2017.07.493>.

### References

Bedgood, D., Boggiano, M.M., Turan, B., 2014. Testosterone and social evaluative stress: the moderating role of basal cortisol. *Psychoneuroendocrinology* 47, 107–115.  
Carré, J.M., Gilchrist, J.D., Morrissey, M.D., McCormick, C.M., 2010. Motivational and situational factors and the relationship between testosterone dynamics and human

aggression during competition. *Biol. Psychol.* 84, 346–353.  
Carré, J.M., Campbell, J.A., Lozoya, E., Goetz, S.M., Welker, K.M., 2013. Changes in testosterone mediate the effect of winning on subsequent aggressive behaviour. *Psychoneuroendocrinology* 38 (10), 2034–2041.  
Edwards, D.A., Casto, K.V., 2015. Baseline cortisol moderates testosterone reactivity to women's intercollegiate athletic competition. *Physiol. Behav.* 142, 48–51.  
Güth, W., Schmittberger, R., Schwarze, B., 1982. An experimental analysis of ultimatum bargaining. *J. Econ. Behav. Organ.* 3, 367–388.  
Galinsky, A.D., Mussweiler, T., 2001. First offers as anchors: the role of perspective-taking and negotiator focus. *J. Pers. Soc. Psychol.* 81 (4), 657–669.  
Geniole, S.N., Bird, B.M., Ruddick, R.L., Carré, J.M., 2017. Effects of competition outcome on testosterone concentrations in humans: an updated meta-analysis. *Horm. Behav.* 92, 37–50.  
Kenny, D.A., Campbell, D.T., 1989. On the measurement of stability in over-time data. *J. Pers.* 57 (2), 445–481. <http://dx.doi.org/10.1111/j.1467-6494.1989.tb00489.x>.  
Kenny, D.A., 1975. Cross-lagged panel correlation: a test for spuriousness. *Psychol. Bull.* 82 (6), 887–903. <http://dx.doi.org/10.1037/0033-2909.82.6.887>.  
Liening, S.H., Stanton, S.J., Saini, E.K., Schultheiss, O.C., 2010. Salivary testosterone, cortisol, and progesterone: two-week stability, interhormone correlations, and effects of time of day, menstrual cycle, and oral contraceptive use on steroid hormone levels. *Physiol. Behav.* 99 (1), 8–16. <http://dx.doi.org/10.1016/j.physbeh.2009.10.001>.  
Maestripieri, D.M., Baran, N., Sapienza, P., Zingales, L., 2010. Between- and within-sex variation in hormonal responses to psychological stress in a large sample of college students. *Stress* 13 (5), 413–424.  
Marceau, K., Ruttle, P.L., Shirtcliff, E.A., Hastings, P.D., Klimes-Dougan, B., Zahn-Waxler, C., 2015. Within-person coupling of changes in cortisol, testosterone, and DHEA across the day in adolescents. *Dev. Psychobiol.* 57 (6), 654–669.  
Mehta, P.H., Josephs, R.A., 2006. Testosterone change after losing predicts the decision to compete again. *Horm. Behav.* 92, 117–127.  
Mehta, P.H., Josephs, R.A., 2010. Testosterone and cortisol jointly regulate dominance: evidence for a dual-hormone hypothesis. *Horm. Behav.* 58 (5), 898–906.  
Mehta, P.H., Mor, S., Yap, A., Prasad, S., 2015a. Dual-hormone changes track bargaining performance. *Psychol. Sci.* 26 (6), 866–876.  
Mehta, P.H., Snyder, N.S., Knight, E.L., Lassetter, B., 2015b. Close versus decisive victory moderates the effect of testosterone change on competitive decisions and task enjoyment. *Adapt. Hum. Behav. Physiol.* 1, 291–311.  
Mehta, P.H., Welker, K.M., Zilioli, S., Carré, J.M., 2015c. Testosterone and cortisol jointly modulate risk-taking. *Psychoneuroendocrinology* 56, 88–99. <http://dx.doi.org/10.1016/j.psyneuen.2015.02.023>.  
Popma, A., Vermeiren, R., Geluk, C.A., Rinne, T., van den Brink, W., Knol, D.L., et al., 2007. Cortisol moderates the relationship between testosterone and aggression in delinquent male adolescents. *Biol. Psychiatry* 61 (3), 405–411.  
Racine, M., Moulin, D.E., Nielson, W.R., Morley-Forster, P.K., Lynch, M., Clark, A.J., et al., 2016. The reciprocal associations between catastrophizing and pain outcomes in patients being treated for neuropathic pain: a cross-lagged panel analysis study. *Pain* 157 (9), 1946–1953. <http://dx.doi.org/10.1097/j.pain.0000000000000594>.  
Rodgers, J.L., Nicewander, W.A., 1988. Thirteen ways to look at the correlation coefficient. *Am. Statistician* 42, 59–66.  
Roest, J.J., Van der Helm, G.H.P., Stams, G.J.J.M., 2016. The relation between therapeutic alliance and treatment motivation in residential youth care: a cross-lagged panel analysis. *Child Adolesc. Soc. Work J.* 33 (5), 455–468. <http://dx.doi.org/10.1007/s10560-016-0438-4>.  
Shirtcliff, E.A., Dismukes, A.R., Marceau, K., Ruttle, P.L., Simmons, J.G., Han, G., 2015. A dual-axis approach to understanding neuroendocrine development. *Dev. Psychobiol.* 57 (6), 643–653.  
Smith, K.M., Apicella, C.L., 2017. Winners, losers, and posers: The effect of power poses on testosterone and risk-taking following competition. *Horm. Behav.* 92, 172–181.  
Spurk, D., Abele, A.E., 2014. Synchronous and time-lagged effects between occupational self-efficacy and objective and subjective career success: findings from a four-wave and 9-year longitudinal study. *J. Vocat. Behav.* 84 (2), 119–132. <http://dx.doi.org/10.1016/j.jvb.2013.12.002>.  
Viau, V., 2002. Functional cross-talk between the hypothalamic-pituitary-gonadal and adrenal axes. *J. Neuroendocrinol.* 14 (6), 506–513.  
Welker, K.M., Lassetter, B., Brandes, C., Prasad, S., Koop, D., Mehta, P.H., 2016. A comparison of salivary testosterone measurement using immunoassays and tandem mass spectrometry. *Psychoneuroendocrinology* 71, 180–188. <http://dx.doi.org/10.1016/j.psyneuen.2016.05.02>.  
Welker, K.M., Norman, R., Goetz, S.M.M., Kitayama, S., Carré, J.M., 2017. Preliminary evidence that testosterone's association with aggressive behavior depends on self-construal. *Horm. Behav.* 92, 117–127.  
Wirtz, C.M., Hofmann, S.G., Riper, H., Berking, M., 2014. Emotion regulation predicts anxiety over a five-year interval: a cross-lagged panel analysis. *Depress. Anxiety* 31 (1), 87–95. <http://dx.doi.org/10.1002/da.22198>.  
Zilioli, S., Watson, N.V., 2013. Winning isn't everything: mood and testosterone regulate the cortisol response in competition. *PLoS One* 8 (1), e52582.